

FALAK PHARMACY
2730 E.TREMONT AVE , BRONX , NY , 10461
Phone#: 7185975230 Fax#: 7185972886

Drug NDC: 10122-0321-05
Drug Name: FACTIVE

How should I use this medicine?

Take this medicine by mouth with a full glass of water. Follow the directions on the prescription label. You can take it with or without food. If it upsets your stomach, take it with food. Take your medicine at regular intervals. Do not take your medicine more often than directed. Take all of your medicine as directed even if you think you are better. Do not skip doses or stop your medicine early. Avoid taking didanosine, sucralfate, antacids, or iron or zinc products within 2 hours of taking this medicine. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.
NOTE: This medicine is only for you. Do not share this medicine with others.

Where should I keep my medicine?

Keep out of the reach of children. Store at room temperature between 15 and 30 degrees C (59 and 86 degrees F). Protect from light. Throw away any unused medicine after the expiration date.

NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.

What side effects may I notice from receiving this medicine?

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash or hives, swelling of the face, lips, or tongue; anxious; breathing problems; bloody or watery diarrhea; confusion; depressed mood; fast, irregular heartbeat; fever; hallucination, loss of contact with reality; joint, muscle, or tendon pain or swelling; loss of memory; pain, tingling, numbness in the hands or feet; seizures; signs and symptoms of high blood sugar such as dizziness; dry mouth; dry skin; fruity breath; nausea; stomach pain; increased hunger or thirst; increased urination; signs and symptoms of liver injury like dark yellow or brown urine; general ill feeling or flu-like symptoms; light-colored stools; loss of appetite; nausea; right upper belly pain; unusually weak or tired; yellowing of the eyes or skin; signs and symptoms of low blood sugar such as feeling anxious; confusion; dizziness; increased hunger; unusually weak or tired; sweating; shakiness; cold; irritable; headache; blurred vision; fast heartbeat; loss of consciousness; pale skin; suicidal thoughts or other mood changes; sunburn; unusually weak or tired. Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): dry mouth; headache; nausea; trouble sleeping.

This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.