

**FALAK PHARMACY**  
**2730 E.TREMONT AVE , BRONX , NY , 10461**  
**Phone#: 7185975230 Fax#: 7185972886**

**Drug NDC: 31722-0562-24**  
**Drug Name: ABACAVIR SULFATE**

**How should I use this medicine?**

Take this medicine by mouth with a glass of water. Follow the directions on the prescription label. Use a specially marked spoon or dropper to measure each dose. Ask your pharmacist if you do not have one. Household spoons are not accurate. You may take this medicine with or without food. Take your medicine at regular intervals. Do not take your medicine more often than directed. For your anti-HIV therapy to work as well as possible, take each dose exactly as prescribed. Do not skip doses or stop your medicine even if you feel better. Skipping doses may make the HIV virus resistant to this medicine and other medicines. Do not stop taking except on your doctor's advice. A special MedGuide will be given to you by the pharmacist with each prescription and refill. Be sure to read this information carefully each time. Talk to your pediatrician regarding the use of this medicine in children. While this drug may be prescribed for children as young as 3 months old for selected conditions, precautions do apply.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once.**  
**NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 20 and 25 degrees C (68 and 77 degrees F). Do not freeze. May be refrigerated. The solution may turn brown over time. Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible: allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue; breathing problems; cough; dizziness; fast, irregular heartbeat; muscle pain or weakness; nausea, vomiting, unusual upset stomach or stomach pain; redness, blistering, peeling or loosening of the skin, including inside the mouth; signs and symptoms of liver injury like dark yellow or brown urine; general ill feeling or flu-like symptoms; light-colored stools; loss of appetite; right upper belly pain; unusually weak or tired; yellowing of the eyes or skin; signs of infection - fever or chills, cough, sore throat, pain or trouble passing urine; Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome): depressed mood; diarrhea; dizziness; headache; trouble sleeping

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**