

**FALAK PHARMACY**  
**2730 E.TREMONT AVE , BRONX , NY , 10461**  
**Phone#: 7185975230 Fax#: 7185972886**

**Drug NDC: 65649-0651-03**

**Drug Name: UCERIS**

**How should I use this medicine?**

This medicine is for rectal use only. Do not take by mouth. Wash hands before and after use. Follow the directions on the prescription label. Before using foam, use the bathroom to empty the bowels. Use foam in a standing, lying or sitting position (e.g., while using the toilet). When using foam in the evening, use right before bedtime and do not try to have a bowel movement again until the next morning. Prepare the canister for use as directed. Use a new applicator for each dose. Place the forefinger on the top of pump dome and then turn the canister upside down. Insert the applicator gently into the rectum as far as it is comfortable. The easiest way to use foam is to keep one foot on the floor and raise the other foot onto a firm surface such as a chair or stool. Use the forefinger to fully push down the pump dome one time and hold it for about 2 seconds in that position. Release finger pressure on the pump dome and hold the applicator in place for 10 to 15 seconds. Remove the applicator. The foam will still expand a little and may drop out of the applicator or anus. Remove the applicator from the canister. Place the used applicator in the plastic bag provided. Throw the plastic bag away in household trash. Do not use your medicine more often than directed. Do not suddenly stop using your medicine because you may develop a severe reaction. Your doctor will tell you how much medicine to use. If your doctor wants you to stop using the medicine, the amount that you use may be slowly lowered over time to avoid any side effects. Talk to your pediatrician regarding the use of this medicine in children. Special care may be needed.

**Overdosage: If you think you have taken too much of this medicine contact a poison control center or emergency room at once. NOTE: This medicine is only for you. Do not share this medicine with others.**

**Where should I keep my medicine?**

Keep out of the reach of children. Store at room temperature between 20 and 25 degrees C (68 and 77 degrees F). Do not refrigerate. The product is flammable. Avoid fire, flames, and smoking during and right after use. Throw away any unused medicine after the expiration date.

**NOTE: This sheet is a summary. It may not cover all possible information. If you have questions about this medicine, talk to your doctor, pharmacist, or health care provider.**

**What side effects may I notice from receiving this medicine?**

Side effects that you should report to your doctor or health care professional as soon as possible:

- allergic reactions like skin rash, itching or hives, swelling of the face, lips, or tongue
- changes in vision
- cough
- fever or chills
- increased thirst
- mental depression, mood swings, mistaken feelings of self-importance, mistaken feelings of being mistreated
- nausea
- pain or trouble passing urine
- skin problems, acne, thin and shiny skin
- sore throat
- swelling of feet or lower legs
- unusual bruising or red pinpoint spots on the skin
- unusually weak or tired
- vomiting
- weight gain
- wounds that will not heal

Side effects that usually do not require medical attention (report to your doctor or health care professional if they continue or are bothersome):

- diarrhea or constipation
- difficulty sleeping
- headache
- increased appetite
- increased sweating
- menstrual problems

**This list may not describe all possible side effects. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**